

Periodicity Schedule

Immunization Schedule

Birth	Hep B
2 months	Pentacel Hep B Pevnar Rotateq
4 months	Pentacel Pevnar Rotateq
6 months	Pentacel Hep B Pevnar Rotateq Flu
9 months	Flu (?)
12 months	MMR Varivax Pevnar Hep A
15 months	Dtap Hib
18 months	Hep A
4 years	Proquad Kinrix
11 years	Menactra Gardasil Tdap
16 years	Menactra Bexsero

Routine Screenings	3-5 day	1 mo	2 mo	4 mo	6 mo	9 mo	12 mo	15 mo	18 mo	24 mo	30 mo	3 yr	4 yr	5 yr	6 yr	7 yr	8 yr	9 yr	10 yr	11 yr	12 yr	13 yr	14 yr	15 yr	16 yr	17 yr	18 yr	
H/H ¹							X																					
Pb							X		X																			
Lipid Panel ²																		X ²	---	---								
BMI									X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	
Blood Pressure												X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	
Hearing												X	X	X			X				X				X		X	
Vision ³												X	X	X	X		X				X				X		X	
Post Partum Depression	X	X	X																									
MCHAT									X	X																		
PEDS (Development)						X			X		X																	
PSC (Mental Health)																		X	X									
Adolescent Questionnaire																					X	X	X	X	X	X	X	
GC/C and HIV ⁴																									X ⁴	---	---	---

Vaccine Trade Names

Pentacel	Dtap, IPV, & Hib
Pevnar	Pneumococcal
Rotateq	Rotavirus
Proquad	MMR & Varicella
Kinrix	Dtap & IPV
Gardasil	HPV
Menactra	Meningococcal ACY W-135
Bexsero	Meningococcal B

Notes

- 1.) **MD:** Assess risk factors for anemia in females 13 yr and older
- 2.) **Clinical Staff/MD:** Check lipid panel once age 9-11
- 3.) **Clinical Staff/MD:** Check vision for all sports physicals
- 4.) **MD:** STI testing: Check GC/C annually if sexually active. Check HIV once age 15-65 (repeat based on risk factors).
- 5.) **Vaccine tips:** Hep A: 2-dose series separated by at least 6 months
Flu: 2 doses separated by at least 1 month 1st yr, then annually
Gardasil: 2-dose series separated by at least 6 months, 9-14 yr
Bexsero: 2-dose series separated by at least 1 month, 16-18 yr (can be given as early as age 10)